

Identifying and Addressing Overbasalization in Outpatient Insulin Therapy Using CGM Data

Nicole E. Siwarski, PharmD; Alex DeLucenay, PharmD, BCACP; Nabila Ahmed-Sarwar, Pharm.D., BCPS, BCACP, CDCES, BC-ADM

Introduction

- The ADA cautions against overbasalization, as it is a risk factor for hypoglycemia; current literature suggests rates of overbasalization range from 5.7-40%¹⁻³
- Overbasalization is defined as a basal insulin dose of ≥ 0.5 units/kg or a bedtime-to-morning (BeAM) difference of ≥ 50 mg/dL^{3,4}
- Continuous glucose monitors (CGM) can be a tool to identify overbasalization in patients and address suboptimal glycemic control⁵

Objectives

- Primary Objective:** Determine the prevalence of insulin overbasalization through CGM utilization
- Secondary Objectives:** Assess changes in glycemic control following CGM initiation and identify patient characteristics associated with overbasalization

Methods

- Retrospective chart review included patients with established care at the internal medicine clinic between January 2023 and November 2025
- Inclusion: Age ≥ 18 years, type 2 diabetes, on basal insulin for ≥ 3 months, at least one HbA1c value within 3 months before and after CGM initiation
- Exclusion: Type 1 diabetes, pregnancy, active malignancy or palliative care, incomplete CGM data ($<70\%$ wear time over 14 days)
- Demographics and clinical data obtained from EMR, CGM data gathered from Dexcom Clarity platform

Results

Baseline Demographics	Entire Population (n=53)	Overbasalized Patients (n=30)
Age (years)	62.2 \pm 10	62.7 \pm 10.7
Female Sex (%)	54.7	62.1
BMI (kg/m ²)	33.7 \pm 9.6	31.8 \pm 6.8
A1c (%)	8.9 \pm 2.1	8.6 \pm 1.7
CGM Data		
Time Very High (%) [*]	8 [IQR 31]	15 [IQR 20.5]
Time High (%)	26 \pm 12.2	27 \pm 9.5
Time in Range (%)	54.3 \pm 27.6	52.8 \pm 22.7
Time Below Range (%)	Negligible (<1)	
Average Glucose (mg/dL)	189.1 \pm 52.6	190.2 \pm 45.3
GMI (%)	7.8 \pm 1.3	7.8 \pm 1.1

* Reported as Median

Average Duration of CGM Use

- 3 months: 105.1 \pm 5.8 days
- 6 months: 197.4 \pm 16.3 days


Time CGM Active

- 3 months: 95.1%
- 6 months: 94.8%


- Overbasalization of insulin was identified in 57% of patients
- CGM data contributed to reduction in overbasalization in 90% patients
- BeAM was reduced from 63.5 mg/dL at baseline to 56.7 mg/dL at 6 months (p=0.09)

ROCHESTER REGIONAL HEALTH | **Rochester General Hospital**


Change in Basal Insulin Dose Over Time



Change in HbA1c Over Time



Change in Time in Range (TIR) Over Time




Corresponding author: nicole.siwarski@rochesterregional.org
ASHP Midyear Clinical Meeting 2025

Results

- 27% of the patients met both the daily basal insulin dose and difference in BeAM criteria for overbasalization.
- Of the overbasalized patients, 67% had follow-up data
- Reduction in overbasalization was observed in 17/20; of those, 8/20 had a decrease in basal insulin dose and 9/20 a decrease in BeAM

Concurrent Diabetes Medication Use at Baseline



- In the overbasalized population:
 - 40% of patients were prescribed prandial insulin at baseline; this decreased to 31% at 3 months, then increased to 44% at 6 months
 - 16.7% were on an atypical antipsychotic and 10% were on a thyroid supplement

Conclusions

- Change in glycemic control based on increased TIR was statistically significant from baseline to 3 months in the overbasalized population
- Change in glycemic control based on decrease in HbA1c in the overbasalized population was not statistically significant
- Hypoglycemia was not observed in the overbasalized population

Discussion

- Prevalence of overbasalization observed higher than that reported in the literature but was not associated with hypoglycemia
- The incidence of overbasalization may be under reported due to follow-up data being available for approximately 2/3 of patients
- The change improvement in glycemic control was observed over time, but reductions in HbA1c and TIR were not sustained overtime
- Thyroid dysfunction likely did not contribute to lack of glycemic control as patients TSH levels were WNL

References

- American Diabetes Association Professional Practice Committee; 7. Diabetes Technology: Standards of Care in Diabetes—2025. *Diabetes Care* 1 January 2025; 48 (Supplement_1): S146-S166. doi:10.2337/dc25-S007.
- Heppen S, Bostan H, Gül Ü, et al. The rate and associated factors of overbasalization in patients with type 2 diabetes mellitus in a tertiary hospital. *Endocrinol Res Pract.* 2025;29(1):3-8. doi:10.5152/ERP.2025.24553.
- Cowart K, Carris NW. Pradlicable measurement and identification of overbasalization. *Clin Diabetes.* 2022;40(1):75-77. doi:10.2337/cd21-0096.
- Stewart-Lynch A, Meyers R, Sidig D, McConville SO, Heiple L. Quantifying and characterizing the presence of insulin overbasalization in a family medicine practice. *Clin Diabetes.* 2024;42(2):266-273. doi:10.2337/cd23-0044.
- Mayberry LS, Guy C, Hendrickson CD, McCoy AB, Elasy TA. Rates and correlates of uptake of continuous glucose monitors among adults with type 2 diabetes in primary care and endocrinology settings. *J Gen Intern Med.* 2023;38(11):2546-2552. doi:10.1007/s11606-023-08222-3

© 2025 ROCHESTER REGIONAL HEALTH. ALL RIGHTS RESERVED. WWW.ROCHESTERREGIONAL.ORG